arXiv:1209.4361v2 [cond-mat.supr-con] 15 Oct 2012

An attempt at a resonating mean-field theoretical description of thermal behavior of two-gap superconductivity

Seiya Nishiyama*, João da Providência[†]
Constança Providência[‡] and Hiromasa Ohnishi[§]

Centro de Física Computacional, Departamento de Física,
Universidade de Coimbra, P-3004-516 Coimbra, Portugal

Nanosystem Research Institute (NRI),
National Institute of Advanced Science and Technology (AIST),
1-1-1, Umezono, Tsukuba, Ibaraki 305-8568, Japan[§]

Dedicated to the Memory of Hideo Fukutome

October 16, 2012

Abstract

The resonating mean-field theory (Res-MFT) has been applied and shown to effectively describe two-gap superconductivity (SC). Particularly at $T = 0$ using a suitable chemical potential, the two-gap SC in MgB_2 has been well described by the Res-Hartree-Bogoliubov theory (Res-HBT). The Res-HB ground state generated with HB wave functions almost exhausts the ground-state correlation energy in all the correlation regimes including an intermediate coupling regime. In this paper we make an attempt at a Res-MF theoretical description of thermal behavior of the two-gap SC. In the single-gap case we find a new formula leading to a higher T_c than the T_c of the usual HB formula.

Keywords: Res-MF theory;BCS model;Two-gap superconductivity

---

*Corresponding author. E-mail address: seikoceu@khe.biglobe.ne.jp

[†] E-mail address: providencia@teor.fis.uc.pt

[‡] E-mail address: cp@teor.fis.uc.pt

[§] E-mail address: hiro.ohnishi@aist.go.jp

1 Introduction

A two-gap superconductivity (SC) of magnesium diboride MgB_2 with critical temperature $T_c = 39$K has been discovered nearly a decade ago [1]. Hitherto, intensive studies had been made to raise the T_c of usual BCS superconductor in the weak coupling regime [2, 3, 4] and to obtain Eliashberg's critical temperature in the strong coupling [5, 6, 7]. The $T_c = 39$K is close to or even above the upper theoretical value predicted by the BCS theory [8]. Through *ab initio* density functional computations it has been estimated as 22K by Kortus et al. [9]. The existence of two energy gaps in MgB_2 has been predicted theoretically by Liu et al. [10] employing the effective σ and π two-band model. They have obtained gaps at $T = 0$, $\Delta_\sigma = 7.4$ [meV] and $\Delta_\pi = 2.4$ [meV] and also their temperature dependencies and $T_c = 40$K. The two-band model was first proposed by Suhl et al. [11] and next introduced by Kondo [12].

To approach such problems, Fukutome has developed the resonating Hartree-Fock theory (Res-HFT) [13] and Fukutome and one of the present authors (S.N.) have extended it directly to the resonating Hartree-Bogoliubov theory (Res-HBT) to include pair correlations [14, 15]. Steadily the Res-HBT has succeeded to describe effectively the two-gap SC [16] (referred to as I). If we get a *Thermal Gap Equation* in the Res-HBT, it is a strong manifestation of analogy of the Res-HBT with the usual BCS and HBT [2, 3, 17, 18]. The Res-HBT has the surprising fact that every HB eigenfunction in an HB resonating state has its own orbital-energy. Due to this fact, the Res-HBT also becomes a possible candidate and is considered to be useful for such subjects. This is because that the Res-HBT has the following characteristic feature: The Res-HBT is equivalent to the coupled Res-HB eigenvalue equations and the orbital concept is still surviving in the Res-HB approximation (Res-HBA) though the orbitals are resonating. This feature permits us to say that in some sense the band picture has a correspondence to the orbital concept in the Res-HBA though bands of different structures are resonating. The Res-HB ground state generated with HB wave functions (WFs) which are the coherent state representations (CS reps) [19], is expected to almost exhausts the ground-state correlation energy in all the correlation regimes including an intermediate coupling regime.

To demonstrate the predominance of the Res-HBT for superconducting fermion systems with large quantum fluctuations over the usual BCS and Eliashberg theories, we already have applied it to a naive BCS Hamiltonian of singlet-pairing. A state with large quantum fluctuations is approximated by superposition of two HB WFs which are non-orthogonal CS reps with different correlation structures. We have optimized directly the Res-MF energy functionals by variations of the Res-MF ground-state energy with respect to the Res-MF parameters, i.e., *energy-gaps.* The Res-MF ground and excited states generated with the two HB WFs explain most of the magnitudes of two energy-gaps in MgB_2. Both the large *energy-gap* and the small one have a significant physical meaning because electron systems, composed of condensed electron pairs, have now strong correlations among the fermions [16, 20].

To go beyond the above *ab initio* density functional computations and phenomenologies, we develop a thermal Res-HBT. A thermal variation is made to satisfy diagonalization conditions for thermal Res-Fock-Bogoliubov (Res-FB) operators along a way very different from the usual thermal-BCS theory [21, 22, 23]. We derive formulas for determining T_c and the behavior of the gap near T=0 and T_c. Particularly in the case of an *equal* magnitude of two gaps but with different phases, we find new formulas for T_c boosting up T_c to higher value than the usual HB's value and get new analytical expressions for gaps near T=0 and T_c. In the intermediate temperature region, we solve numerically a thermal resonating gap equation.

2 Thermal resonating HB eigenvalue equation

Consider the whole Res-HB subspace $|\Psi^{\mathrm{Res}(k)}\rangle = \sum_{r=1}^{n} c_r^{(k)} |g_r\rangle (|g_r\rangle;\mathrm{HB\ WFs})$, $(k=1,\cdots,n)$, in which the Res-state with index $k=1$ and the Res-states with indices $k=2,\cdots,n$ stand for the Res-ground state and the Res-excited states, respectively. We introduce a temperature dependent Lagrangian with the Lagrange multiplier term $E^{(k)}$ to secure the normalization condition $\langle\Psi^{\mathrm{Res}(k)}|\Psi^{\mathrm{Res}(k)}\rangle = 1$:

$$L[\mathcal{Z}]_{\mathrm{Res}}^{\mathrm{HB}} = \sum_{k=1}^{n} \sum_{r,s=1}^{n} \left\{ H\left[W_{rs}[\mathcal{Z}]_{\mathrm{Res}}\right] - E^{(k)} \right\} \cdot [\det z_{rs}]^{\frac{1}{2}}\, c_r^{(k)*} c_s^{(k)},\ \ \mathcal{Z} = \sum_{r=1}^{n} \oplus \mathcal{F}_r, \tag{2.1}$$

variation of which is made in a quite parallel manner to the one in the previous works where $\mathcal{F}_r (r=1,\cdots,n)$ is a Res-FB operator [13, 14]. From the variation of $L[\mathcal{Z}]_{\mathrm{Res}}^{\mathrm{HB}}$ with respect to $c_r^{(k)*}$ for any k, we get a thermal Res-HB CI equation to determine $c_s^{(k)}$

$$\sum_{s=1}^{n} \left\{ H\left[W_{rs}[\mathcal{Z}]_{\mathrm{Res}}\right] - E^{(k)} \right\} \cdot [\det z_{rs}]^{\frac{1}{2}} c_s^{(k)} = 0. \tag{2.2}$$

We define D_{rs} and $\tilde{D}_{rs}$ as $D_{rs} \equiv u_s z_{rs}^{-1} \delta u_r^{\dagger}$ and $\tilde{D}_{rs} \equiv \delta u_s z_{rs}^{-1} u_r^{\dagger}$, respectively, where $z_{rs} \equiv u_r^{\dagger} u_s$ and $u_r^{\mathrm{T}} \equiv [b_r^{\mathrm{T}}, a_r^{\mathrm{T}}]$ (a_r and b_r:r-th HB amplitudes). The variations of thermal HB interstate density matrix $W[\mathcal{Z}]_{\mathrm{Res}}$ and thermal overlap integral $[\det z]^{1/2}$ are given by

$$\left.\begin{aligned} \delta W_{rs}[\mathcal{Z}]_{\mathrm{Res}} &= D_{rs}(1_{2N} - W_{rs}[\mathcal{Z}]_{\mathrm{Res}}) + (1_{2N} - W_{rs}[\mathcal{Z}]_{\mathrm{Res}})\tilde{D}_{rs}, \\ \delta[\det z_{rs}]^{\frac{1}{2}} &= \frac{1}{2}\mathrm{Tr}(D_{rs} + \tilde{D}_{rs}) \cdot [\det z_{rs}]^{\frac{1}{2}}. \end{aligned}\right\} \tag{2.3}$$

We obtain also the variation of Hamiltonian matrix element $H[W_{rs}[\mathcal{Z}]_{\mathrm{Res}}]$ as

$$\left.\begin{aligned} \delta H[W_{rs}[\mathcal{Z}]_{\mathrm{Res}}] &= \frac{1}{2}\mathrm{Tr}\{\mathcal{F}[W_{rs}[\mathcal{Z}]_{\mathrm{Res}}]\delta W_{rs}[\mathcal{Z}]_{\mathrm{Res}}\}, \\ \mathcal{F}[W_{rs}[\mathcal{Z}]_{\mathrm{Res}}] &= \begin{bmatrix} F_{rs}[\mathcal{Z}]_{\mathrm{Res}} & D_{rs}[\mathcal{Z}]_{\mathrm{Res}} \\ -D_{sr}^{*}[\mathcal{Z}]_{\mathrm{Res}} & -F_{sr}^{*}[\mathcal{Z}]_{\mathrm{Res}} \end{bmatrix}, \\ F_{rs;\alpha\beta}[\mathcal{Z}]_{\mathrm{Res}} &\equiv \frac{\delta H[W_{rs}[\mathcal{Z}]_{\mathrm{Res}}]}{\delta R_{rs;\beta\alpha}[\mathcal{Z}]_{\mathrm{Res}}} = h_{\alpha\beta} + [\alpha\beta|\gamma\delta] R_{rs;\delta\gamma}[\mathcal{Z}]_{\mathrm{Res}}, \\ D_{rs;\alpha\beta}[\mathcal{Z}]_{\mathrm{Res}} &\equiv \frac{\delta H[W_{rs}[\mathcal{Z}]_{\mathrm{Res}}]}{\delta K_{sr;\alpha\beta}^{*}[\mathcal{Z}]_{\mathrm{Res}}} = -\frac{1}{2}[\alpha\gamma|\beta\delta] K_{rs;\delta\gamma}[\mathcal{Z}]_{\mathrm{Res}}. \end{aligned}\right\} \tag{2.4}$$

Following [14] and the Res-HF theory [13], writing $L[\mathcal{Z}]_{\mathrm{Res}}^{\mathrm{HB}} = \sum_{k=1}^{n} \sum_{r,s=1}^{n} \mathcal{L}_{rs}^{\mathrm{HB(k)}}[\mathcal{Z}]_{\mathrm{Res}} c_r^{(k)*} c_s^{(k)}$ and $\mathcal{L}_{rs}^{\mathrm{HB(k)}}[\mathcal{Z}]_{\mathrm{Res}} = \{H[W_{rs}[\mathcal{Z}]_{\mathrm{Res}}] - E^{(k)}\} \cdot [\det z_{rs}]^{1/2}$, from the variation of $L_{\mathrm{Res}}^{\mathrm{HB}}$ for any k, we obtain a thermal Res-HB equation to determine the thermal MF WF u_r as

$$\left.\begin{aligned} \sum_{k=1}^{n} \sum_{s=1}^{n} \mathcal{K}_{rs}^{(k)}[\mathcal{Z}]_{\mathrm{Res}} c_r^{(k)*} c_s^{(k)} &= 0, \\ \mathcal{K}_{rs}^{(k)}[\mathcal{Z}]_{\mathrm{Res}} \equiv \left\{ (1_{2N} - W_{rs}[\mathcal{Z}]_{\mathrm{Res}}) \mathcal{F}[W_{rs}[\mathcal{Z}]_{\mathrm{Res}}] + H[W_{rs}[\mathcal{Z}]_{\mathrm{Res}}] - E^{(k)} \right\} &\cdot W_{rs}[\mathcal{Z}]_{\mathrm{Res}} \cdot [\det z_{rs}]^{\frac{1}{2}}. \end{aligned}\right\} \tag{2.5}$$

From (2.5) we can derive a thermal Res-HB eigenvalue equation which is expressed as follows:

$$\left.\begin{aligned} &[\mathcal{F}_r[\mathcal{Z}]_{\mathrm{Res}} u_r]_i = \epsilon_{ri} u_{ri}, \quad \epsilon_{ri} \equiv \widetilde{\epsilon}_{rs} - \sum_{k=1}^{n} \left\{ H[W_{rr}[\mathcal{Z}]_{\mathrm{Res}}] - E^{(k)} \right\} |c_r^{(k)}|^2, \\ &\mathcal{F}_r[\mathcal{Z}]_{\mathrm{Res}} = \mathcal{F}_r[\mathcal{Z}]_{\mathrm{Res}}^{\dagger} \equiv \mathcal{F}[W_{rr}[\mathcal{Z}]_{\mathrm{Res}}] \sum_{k=1}^{n} |c_r^{(k)}|^2 \\ &\qquad\qquad + \sum_{k=1}^{n} {\sum_{s=1}^{\prime\, n}} \left\{ \mathcal{K}_{rs}^{(k)}[\mathcal{Z}]_{\mathrm{Res}} c_r^{(k)*} c_s^{(k)} + \mathcal{K}_{rs}^{(k)\dagger}[\mathcal{Z}]_{\mathrm{Res}} c_r^{(k)} c_s^{(k)*} \right\}. \end{aligned}\right\} \tag{2.6}$$

We call $\mathcal{F}_r[\mathcal{Z}]_{\mathrm{Res}}$ the thermal Res-FB operator. For later convenience we denote $W_{rs}[\mathcal{Z}]_{\mathrm{Res}}$ and $\mathcal{F}_r[\mathcal{Z}]_{\mathrm{Res}}$ simply as W_{rs} and $\mathcal{F}_r$. The Res-HB eigenvalue equation $\mathcal{F}_r u_{rp} = \epsilon_{rp} u_{rp}$ obtained in [14], thus, can be extended to the thermal Res-HB eigenvalue equation (2.6) in a formal way.

3 Thermal resonating gap equation

The thermal HB density matrix is given as $W_{rrp}[\mathcal{F}_{rp}] = [1 + \exp\{\beta\mathcal{F}_{rp}\}]^{-1} (\beta = 1/k_B T, r = 1, 2)$ in momentum p. Using a Bogoliubov transformation $g_{1(2)p}$, $W_{11(22)p}[\mathcal{F}_{1(2)p}]$ are diagonalized as

$$\widetilde{W}_{rp} = g^\dagger_{rp} W_{rrp}[\mathcal{F}_{rp}] g_{rp} = \begin{bmatrix} \widetilde{w}_{rp} & 0 \\ 0 & 1 - \widetilde{w}_{rp} \end{bmatrix}, \ \widetilde{w}_{rp} = \frac{1}{1 + e^{\beta\widetilde{\epsilon}_{rp}}}, \ 1 - \widetilde{w}_{rp} = \frac{1}{1 + e^{-\beta\widetilde{\epsilon}_{rp}}}, \tag{3.1}$$

which are the generalizations of the usual results to the Res-MFT. This is proved elsewhere. By making the Bogoliubov transformation g_{rp}, eigenvalues $\widetilde{\epsilon}_{rp}$ are obtained by diagonalization of the thermal Res-FB operators $\mathcal{F}_{rp}$ with additional terms $(H[W_{rrp}] - E)|c_{rp}|^2$ $(r = 1, 2)$. The thermal HB interstate density matrix in the whole Res-HB subspace is given as the direct sum:

$$W_p[\mathcal{F}_p] = g_p \widetilde{W}_p g^\dagger_p = \bigoplus\nolimits_{r=1}^{2} W_{rrp}[\mathcal{F}_{rp}], \quad W_{rr}[\mathcal{F}_r] = g_{rp}\widetilde{W}_{rp} g^\dagger_{rp}. \tag{3.2}$$

Suppose a *tilde* thermal Res-HB density operator $\widetilde{W}_{1(2)p}$ for *equal-gaps* to be

$$\widetilde{W}_{1(2)p} = \begin{bmatrix} \widetilde{W}^{\uparrow}_{1(2)p} \cdot I_2 & 0 \\ 0 & \widetilde{W}^{\downarrow}_{1(2)p} \cdot I_2 \end{bmatrix}, \ \widetilde{W}^{\uparrow(\downarrow)}_{1(2)p} = \begin{bmatrix} \widetilde{w}^{\uparrow(\downarrow)}_{1(2)p} \cdot I_2 & 0 \\ 0 & (1 - \widetilde{w}^{\uparrow(\downarrow)}_{1(2)p}) \cdot I_2 \end{bmatrix}. \tag{3.3}$$

Here I_2 is the two-dimensional unit matrix. Performing the unitary transformation by $\widehat{g}^{\uparrow(\downarrow)}_{1(2)p}$, we obtain the following thermal Res-HB density matrix $\widetilde{W}^{\uparrow(\downarrow)}_{1(2)p}$.

$$\begin{aligned} W^{\uparrow(\downarrow)}_{1(2)p} &= \widehat{g}^{\uparrow(\downarrow)}_{1(2)p} \widetilde{W}^{\uparrow(\downarrow)}_{1(2)p} \widehat{g}^{\uparrow(\downarrow)\dagger}_{1(2)p} = \widehat{g}^{\uparrow(\downarrow)}_{1(2)p} \begin{bmatrix} \widetilde{w}^{\uparrow(\downarrow)}_{1(2)p} \cdot I_2 & 0 \\ 0 & (1 - \widetilde{w}^{\uparrow(\downarrow)}_{1(2)p}) \cdot I_2 \end{bmatrix} \widehat{g}^{\uparrow(\downarrow)\dagger}_{1(2)p} \\ &= \begin{bmatrix} \frac{1}{2}\left\{1 - \cos\widehat{\theta}_{1(2)p}\left(1 - 2\widetilde{w}^{\uparrow(\downarrow)}_{1(2)p}\right)\right\} \cdot I_2 & \{-(+)\} \times \frac{1}{2}\sin\widehat{\theta}_{1(2)p} e^{-i\widehat{\psi}_{1(2)}}\left(1 - 2\widetilde{w}^{\uparrow(\downarrow)}_{1(2)p}\right) \cdot I_2 \\ \{-(+)\} \times \frac{1}{2}\sin\widehat{\theta}_{1(2)p} e^{i\widehat{\psi}_{1(2)}}\left(1 - 2\widetilde{w}^{\uparrow(\downarrow)}_{1(2)p}\right) \cdot I_2 & \frac{1}{2}\left\{1 + \cos\widehat{\theta}_{1(2)p}\left(1 - 2\widetilde{w}^{\uparrow(\downarrow)}_{1(2)p}\right)\right\} \cdot I_2 \end{bmatrix}. \end{aligned} \tag{3.4}$$

In the *equal-gaps* case $(H[W_{11}] = H[W_{22}] = H[W])$, following I, the Res-HB ground (excited) energy $E^{\mathrm{Res}}_{\mathrm{gr(ex)}}$ is classified into two cases, according to the solutions for the Res-HB CI equation:

$$\left.\begin{aligned} \text{Case I} \quad &: \quad H[W] - H[W_{12}] > 0, \\ E^{\mathrm{Res}}_{\mathrm{gr(ex)}} \ &= \ \frac{1}{1 + (-)[\det z_{12}]^{\frac{1}{2}}} \cdot \left(H[W] + (-)H[W_{12}] \cdot [\det z_{12}]^{\frac{1}{2}}\right), \end{aligned}\right\} \tag{3.5}$$

$$\left.\begin{aligned} \text{Case II} \quad &: \quad H[W] - H[W_{12}] < 0, \\ E^{\mathrm{Res}}_{\mathrm{gr(ex)}} \ &= \ \frac{1}{1 - (+)[\det z_{12}]^{\frac{1}{2}}} \cdot \left(H[W] - (+)H[W_{12}] \cdot [\det z_{12}]^{\frac{1}{2}}\right). \end{aligned}\right\} \tag{3.6}$$

The Res-FB operator $\mathcal{F}^{\uparrow}_{1(2)p}$ for spin-up state, accompanying quantities with upper or lower sign corresponding to Case I (3.5) and Case II (3.6), is expressed as

$$\mathcal{F}^{\uparrow}_{1(2)p} = \begin{bmatrix} \mathcal{F}^{\uparrow}_{+\varepsilon_p} \cdot I_2 & \{+(-)\} \times \mathcal{F}^{\uparrow}_{\Delta p} \cdot I_2 \\ \{+(-)\} \times \mathcal{F}^{\uparrow}_{\Delta p} \cdot I_2 & -\mathcal{F}^{\uparrow}_{-\varepsilon_p} \cdot I_2 \end{bmatrix}. \tag{3.7}$$

The quantities $\mathcal{F}^{\uparrow}_{+\varepsilon_p}, \mathcal{F}^{\uparrow}_{-\varepsilon_p}$ and $\mathcal{F}^{\uparrow}_{\Delta p}$ for Case I(upper sign) and Case II(lower sign) are defined as

$$\mathcal{F}^{\uparrow}_{+(-)\varepsilon_p} \equiv \frac{1}{2}\left\{\varepsilon_p + 2\widetilde{E}^{\mathrm{Res}}_{\mathrm{gr}} \frac{\sin^2\frac{\theta_p}{2}\left(\cos^2\frac{\theta_p}{2}\right)}{\cos\theta_p} \mp \frac{\Delta^2}{\varepsilon_p} \cdot [\det z_{12}]^{\frac{1}{2}}\right\} \cdot \frac{1}{1 \pm [\det z_{12}]^{\frac{1}{2}}}, \widetilde{E}^{\mathrm{Res}}_{\mathrm{gr}} \equiv H[W] - E^{\mathrm{Res}}_{\mathrm{gr}}, \tag{3.8}$$

$$\mathcal{F}^{\uparrow}_{\Delta p}=\mathcal{F}^{\uparrow}_{\Delta}\equiv-\frac{1}{2}\Delta\left\{N(0)V\cdot\mathrm{arcsinh}\left(\frac{1}{x}\right)\pm[\det z_{12}]^{\frac{1}{2}}\right\}\cdot\frac{1}{1\pm[\det z_{12}]^{\frac{1}{2}}},\ \left(x=\frac{\Delta}{\hbar\omega_D}\right), \tag{3.9}$$

where

$$[\det z_{12}]^{\frac{1}{2}}=\exp\left[-2N(0)\hbar\omega_D\left\{\ln(1+x^2)+2x\cdot\arctan\left(\frac{1}{x}\right)\right\}\right]. \tag{3.10}$$

At finite temperature, the quantities $\mathcal{F}^{\uparrow}_{+\varepsilon_p}$, $\mathcal{F}^{\uparrow}_{-\varepsilon_p}$, $\mathcal{F}^{\uparrow}_{\Delta p}$ and Δ become temperature-dependent. This is explicitly expressed by a subscript T. Using the distributions (3.1) we require correspondence relations $\cos\theta_{pT}\Rightarrow\cos\widehat{\theta}_{1p}$ and $\sin\theta_{pT}\Rightarrow\sin\widehat{\theta}_{1p}$ given through

$$\left.\begin{aligned}\cos\theta_p&=\frac{\varepsilon_p}{\sqrt{\varepsilon_p^2+\Delta_T^2}}=\frac{\mathcal{F}^{\uparrow}_{+\varepsilon_pT}+\mathcal{F}^{\uparrow}_{-\varepsilon_pT}}{2\,\widetilde{\epsilon}^{\uparrow}_{1p}}\left(1-2\widetilde{w}^{\uparrow}_{1p}\right),\\ \sin\theta_p&=\frac{\Delta_T}{\sqrt{\varepsilon_p^2+\Delta_T^2}}=-\frac{\mathcal{F}^{\uparrow}_{\Delta_T}}{\widetilde{\epsilon}^{\uparrow}_{1p}}\left(1-2\widetilde{w}^{\uparrow}_{1p}\right),\end{aligned}\right\}\rightarrow\frac{\Delta_T}{\varepsilon_p}=-\frac{\mathcal{F}^{\uparrow}_{\Delta_T}\left(1-2\widetilde{w}^{\uparrow}_{1p}\right)}{\frac{\mathcal{F}^{\uparrow}_{+\varepsilon_pT}+\mathcal{F}^{\uparrow}_{-\varepsilon_pT}}{2}\left(1-2\widetilde{w}^{\uparrow}_{1p}\right)}. \tag{3.11}$$

Notice the multiplication factor $1-2\widetilde{w}^{\uparrow(\downarrow)}_{1p}$. The $\widetilde{\epsilon}^{\uparrow}_{1p}(=\widetilde{\epsilon}^{\uparrow}_{p})$ is the quasi-particle (QP) energy: $\widetilde{\epsilon}^{\uparrow}_{p}=\{(\mathcal{F}^{\uparrow}_{+\varepsilon_pT}+\mathcal{F}^{\uparrow}_{-\varepsilon_pT})^2/4+\mathcal{F}^{\uparrow 2}_{\Delta_T}\}^{1/2}$. The first two equations in parenthesis of (3.11) are unified into the second single equation in (3.11). From now it is shown that equation (3.11) plays the role of the self-consistency condition at finite temperature. Dividing numerator and denominator, respectively by $(\varepsilon_p^2+\Delta_T^2)^{3/2}$ in R.H.S. of the second equation in (3.11), we have

$$1=\frac{\frac{\varepsilon_p^2}{(\varepsilon_p^2+\Delta_T^2)^{\frac{3}{2}}}\left(-\frac{2\mathcal{F}^{\uparrow}_{\Delta_T}}{\Delta_T}\right)\left(1-2\widetilde{w}^{\uparrow}_{p}\right)}{\frac{\varepsilon_p}{(\varepsilon_p^2+\Delta_T^2)^{\frac{3}{2}}}\left(\mathcal{F}^{\uparrow}_{+\varepsilon_pT}+\mathcal{F}^{\uparrow}_{-\varepsilon_pT}\right)\left(1-2\widetilde{w}^{\uparrow}_{p}\right)},\quad \widetilde{w}^{\uparrow}_{p}=\widetilde{w}^{\uparrow}_{1p}. \tag{3.12}$$

Now we demand a new condition for *Thermal Gap Equation*

$$\textstyle\sum_p\left\{\frac{\varepsilon_p}{(\varepsilon_p^2+\Delta_T^2)^{\frac{3}{2}}}\frac{1}{2}\left(\mathcal{F}^{\uparrow}_{+\varepsilon_pT}+\mathcal{F}^{\uparrow}_{-\varepsilon_pT}\right)-\frac{\varepsilon_p^2}{(\varepsilon_p^2+\Delta_T^2)^{\frac{3}{2}}}\left(-\frac{\mathcal{F}^{\uparrow}_{\Delta_T}}{\Delta_T}\right)\right\}\left(1-2\widetilde{w}^{\uparrow}_{p}\right)=0, \tag{3.13}$$

which leads to

$$\left.\begin{aligned}&\left\{1-N(0)V\cdot\mathrm{arcsinh}\left(\frac{1}{x_T}\right)\mp[\det z_{12}]^{\frac{1}{2}}_T\right\}\textstyle\sum_p A_p\\ &\qquad\qquad+\widetilde{E}^{\mathrm{Res}(\pm)}_{\mathrm{gr}_T}\hbar\omega_D\textstyle\sum_p B_p\mp\Delta_T^2\cdot[\det z_{12}]^{\frac{1}{2}}_T\textstyle\sum_p C_p=0,\\ &\frac{\widetilde{E}^{\mathrm{Res}(\pm)}_{\mathrm{gr}_T}}{\hbar\omega_D}=\pm 2N(0)\hbar\omega_D x_T^2\cdot\mathrm{arcsinh}\left(\frac{1}{x_T}\right)\left\{2-N(0)V\cdot\mathrm{arcsinh}\left(\frac{1}{x_T}\right)\right\}\frac{[\det z_{12}]^{\frac{1}{2}}_T}{1\pm[\det z_{12}]^{\frac{1}{2}}_T},\end{aligned}\right\} \tag{3.14}$$

which we have calculated using the solutions for the Res-HB CI equation obtained in I. We also give the following definitions for $\sum_p A_p$, $\sum_p B_p$ and $\sum_p C_p$:

$$\left[\textstyle\sum_p A_p,\ \sum_p B_p,\ \sum_p C_p\right]\equiv\textstyle\sum_p\left[\frac{\varepsilon_p^2}{(\varepsilon_p^2+\Delta_T^2)^{\frac{3}{2}}},\ \frac{1}{\varepsilon_p^2+\Delta_T^2},\ \frac{1}{(\varepsilon_p^2+\Delta_T^2)^{\frac{3}{2}}}\right]\left(1-2\widetilde{w}^{\uparrow}_{p}\right). \tag{3.15}$$

Rearranging (3.14), it is cast to a *Thermal Gap Equation* with similar form to the one in I:

$$\begin{aligned}\frac{1}{N(0)V}&=\mathrm{arcsinh}\left(\frac{1}{x_T}\right)\left[1\pm 2N(0)\frac{\mathrm{arcsinh}\left(\frac{1}{x_T}\right)}{\sum_p A_p}\frac{\Delta_T\cdot\Delta_T\sum_p B_p}{1\pm[\det z_{12}]^{\frac{1}{2}}_T}\cdot[\det z_{12}]^{\frac{1}{2}}_T\right]\\ &\times\left[1+\left\{\mp 1\mp\frac{\Delta_T^2\sum_p C_p}{\sum_p A_p}\pm 4N(0)\frac{\mathrm{arcsinh}\left(\frac{1}{x_T}\right)}{\sum_p A_p}\frac{\Delta_T\cdot\Delta_T\sum_p B_p}{1\pm[\det z_{12}]^{\frac{1}{2}}_T}\right\}\cdot[\det z_{12}]^{\frac{1}{2}}_T\right]^{-1}.\end{aligned} \tag{3.16}$$

The summations $\sum_p A_p$, $\sum_p B_p$ and $\sum_p C_p$ near $T=0$ are computed in Appendix A. Substituting (A.1), (A.12) and (A.11) into (3.16), then, near $T=0$ we have the gaps for Case I (3.5) as

$$\Delta_T^{\mathrm{I}} = \Delta_0 \left[1 - \frac{[\det z_{12}]_0^{\frac{1}{2}}}{N(0)V} \left\{ \operatorname{arcsinh}\left(\frac{\hbar\omega_D}{\Delta_0}\right) - 1 + \frac{[\det z_{12}]_0^{\frac{1}{2}}}{N(0)V} \right\}^{-1} \cdot T_{\frac{1}{2}}^{\mathrm{I}} \right]. \tag{3.17}$$

On the other hand, with the aid of $[\det z_{12}]_0^{1/2} \simeq 1-2\pi N(0)\hbar\omega_D x_0 (0<x_0\ll 1)$ which is easily derived from Taylor expansion of (3.10), we obtain the gap for Case II (3.6) as

$$\Delta_T^{\mathrm{II}} = \Delta_0 \left[1+\frac{1}{\pi\sqrt{1+(N(0)V)^2}} \left\{ \pi-\left(1-N(0)V-\sqrt{1+(N(0)V)^2}\right) \operatorname{arcsinh}\left(\frac{\hbar\omega_D}{\Delta_0}\right) \right\} \cdot T_{\frac{1}{2}}^{\mathrm{II}} \right], \tag{3.18}$$

where we have used the approximate relation

$$\frac{1}{2} N(0)V \operatorname{arcsinh}\left(\frac{\hbar\omega_D}{\Delta_0}\right) \left\{ \operatorname{arcsinh}\left(\frac{\hbar\omega_D}{\Delta_0}\right) -1-1 \right\} \approx \operatorname{arcsinh}\left(\frac{\hbar\omega_D}{\Delta_0}\right) -1. \tag{3.19}$$

In (3.17) and (3.18), as shown from Appendix A, the quantity $T_{\frac{n}{2}}^{\mathrm{I(II)}}$ is defined by

$$T_{\frac{n}{2}}^{\mathrm{I(II)}} \equiv \sqrt{2\pi} \left\{ \widetilde{\Delta}_T^{\mathrm{I(II)}} \right\}^{-\frac{n}{2}} \left\{ \frac{k_B T}{\hbar\omega_D} \right\}^{\frac{n}{2}} \exp\left(-\frac{\hbar\omega_D}{k_B T} \widetilde{\Delta}_T^{\mathrm{I(II)}} \right). \ (n=1,3,\cdots) \tag{3.20}$$

In the opposite limit $T \to T_c^{\mathrm{I}}$ (T_c for Case I) the gap becomes very small, $[\det z_{12}]_T^{1/2} \to 1$, then $\mathcal{F}_{\Delta_T}^{\uparrow} \to -\Delta N(0)V \operatorname{arcsinh}(\hbar\omega_D/\Delta)/4$ and $(\mathcal{F}_{+\varepsilon_p T}^{\uparrow} + \mathcal{F}_{-\varepsilon_p T}^{\uparrow})/2 \to \varepsilon_p/4$ if we use the relations (3.8) and (3.9). We have an approximate QP energy $\widetilde{\epsilon}_p^{\uparrow\mathrm{I}} \simeq [\varepsilon_p^2 + \{-\Delta N(0)V \operatorname{arcsinh}(\hbar\omega_D/\Delta)\}^2]^{1/2}/4$ in which the appearance of numerical factor 1/4 should be become aware carefully. This is because two HB WFs have different correlation structures $\psi_2=\pi$ and $\psi_1=0$. In such a case, returning to the original form of the BCS gap equation but with the modified QP energy $\widetilde{\epsilon}_p^{\uparrow\mathrm{I}}$, the thermal gap equation is expressed as $1=V/2\sum_p \left(1-2\widetilde{w}_p^{\uparrow}\right)/\widetilde{\epsilon}_p^{\uparrow\mathrm{I}}$ and leads to the integral form

$$1=4N(0)V\int_0^{\hbar\omega_D} d\varepsilon \frac{1}{\varepsilon} \tanh\left(\frac{\varepsilon}{8k_B T^{\mathrm{I}}}\right). \tag{3.21}$$

We should emphasize that this form results from the above numerical factor 1/4. Introduce a dimensionless variable $y_T^{\mathrm{I}} \equiv \varepsilon/8k_B T^{\mathrm{I}}$, its upper-value $y_{T_c^{\mathrm{I}}}^{\mathrm{I}} \equiv \hbar\omega_D/8k_B T_c^{\mathrm{I}}$ and the Debye temperature $\theta_D \equiv \hbar\omega_D/k_B$. Integrating R.H.S. of (3.21) by parts, it is approximated as follows:

$$\frac{1}{4N(0)V} \simeq \ln y_{T_c^{\mathrm{I}}}^{\mathrm{I}} - \int_0^{\infty} dy \ln y \ \mathrm{sech}^2 y = \ln y_{T_c^{\mathrm{I}}}^{\mathrm{I}} + \ln\left(\frac{4e^C}{\pi}\right) = \ln\left(\frac{e^C}{2\pi}\frac{\hbar\omega_D}{k_B T_c^{\mathrm{I}}}\right) \equiv \ln\left(\frac{\theta_D}{\widetilde{T}_c^{\mathrm{I}}}\right), \tag{3.22}$$

where we have used the formula given in textbook [24]. The number C is the Euler's constant ($C\simeq 0.5772$) and $e^C \simeq 1.781$. Finally a small rearrangement yields

$$T_c^{\mathrm{I}} = 0.283\theta_D e^{-\frac{1}{4N(0)V}}. \tag{3.23}$$

It should be compared with the Eliashberg's formula [25] and the usual HB's one for T_c [8]

$$T_c = 1.130\theta_D e^{-\frac{1}{N(0)V}}. \tag{3.24}$$

The new formula (3.23) gives a *high* critical-temperature, e.g., T_c^{I}=72.87K for $N(0)V$=0.25 and θ_D=700K. This T_c^{I} is in contrast to T_c obtained by the famous HB formula (3.24) given by Rickayzen and Cohen in [8], namely, T_c=14.49K for the same values of $N(0)V$ and θ_D.

From now we discuss behaviour of the gap near T_c. In the above the modified QP energy $\widetilde{\epsilon}=[\varepsilon^2+\{-\Delta N(0)V \operatorname{arcsinh}(\hbar\omega_D/\Delta)\}^2]^{1/2}/4$ plays a crucial role to boost the T_c (3.23) comparing with T_c=14.49K by (3.24). Note the numerical factor 1/4 in $\widetilde{\epsilon}$. First consider Δ_T^{I} near T_c^{I}. Using this form of the modified QP energy, the gap equation is roughly rewritten as

$$\begin{aligned} \frac{1}{4N(0)V} &= \frac{1}{4}\int_0^{\hbar\omega_D} d\varepsilon \frac{1}{\widetilde{\epsilon}} \tanh\left(\frac{\widetilde{\epsilon}}{2k_B T}\right) \simeq \int_0^{\hbar\omega_D} d\varepsilon \frac{1}{\varepsilon} \tanh\left(\frac{\varepsilon}{8k_B T}\right) \\ &- \left\{ \Delta_T N(0)V \cdot \operatorname{arcsinh}\left(\frac{\hbar\omega_D}{\Delta_T}\right) \right\}^2 \int_0^{\hbar\omega_D} d\varepsilon \left\{ \frac{1}{\varepsilon^3} \tanh\left(\frac{\varepsilon}{8k_B T}\right) - \frac{1}{\varepsilon^2}\frac{1}{8k_B T} \mathrm{sech}^2\left(\frac{\varepsilon}{8k_B T}\right) \right\}, \end{aligned} \tag{3.25}$$

from which we obtain

$$\frac{1}{4N(0)V}=\ln\left(\frac{\hbar\omega_D}{k_B\widetilde{T}}\right)-\frac{7}{8\pi^2}\zeta(3)\left(\frac{2\pi}{e^C}\right)^2\left(\frac{\hbar\omega_D}{k_B\widetilde{T}}\right)^2\left\{N(0)Vx_T\cdot\mathrm{arcsinh}\left(\frac{1}{x_T}\right)\right\}^2$$
$$\simeq\ln\left(\frac{\hbar\omega_D}{k_B\widetilde{T}_c^{\mathrm{I}}}\right)+\frac{\widetilde{T}_c^{\mathrm{I}}-\widetilde{T}}{\widetilde{T}_c^{\mathrm{I}}}-\frac{7}{8\pi^2}\zeta(3)\left(\frac{2\pi}{e^C}\right)^2\left(1-\frac{\widetilde{T}_c^{\mathrm{I}}-\widetilde{T}}{\widetilde{T}_c^{\mathrm{I}}}\right)^{-2}\left(\frac{\hbar\omega_D}{k_B\widetilde{T}_c^{\mathrm{I}}}\right)^2\left\{N(0)Vx_T\cdot\mathrm{arcsinh}\left(\frac{1}{x_T}\right)\right\}^2, \tag{3.26}$$

where $\hbar\omega_D/k_B\widetilde{T}\equiv e^C/2\pi\cdot\hbar\omega_D/k_BT$ and we have used the famous integral-formula (A.18). Using $\mathrm{arcsinh}(1/x_T)\simeq\ln(2/x_T)$ $(0<x_T<1)$, (3.22) and (3.26), we get Δ_T^{I} near T_c^{I} as

$$\Delta_T^{\mathrm{I}}\simeq 2\pi\sqrt{\frac{2}{7\zeta(3)}\frac{k_BT_c^{\mathrm{I}}}{N(0)V}}\left(1-\frac{T_c^{\mathrm{I}}-T}{T_c^{\mathrm{I}}}\right)\sqrt{\frac{T_c^{\mathrm{I}}-T}{T_c^{\mathrm{I}}}}. \tag{3.27}$$

Such a formula has been brought through the use of the modified QP energy $\widetilde{\epsilon}$ which owes to a resonant feature of the multi-band SC. This new formula shows a more complicated temperature-dependence of Δ_T^{I} than the $\sqrt{T_c^{\mathrm{I}}}\sqrt{T_c^{\mathrm{I}}-T}$ dependence of Δ_T^{I} presented by equation (36.6) in textbook [22] and by equations (16.32) and (16.33) in textbook [23], respectively.

Next, for Case II, $(\mathcal{F}^{\uparrow}_{+\varepsilon_pT}+\mathcal{F}^{\uparrow}_{-\varepsilon_pT})$, $\mathcal{F}^{\uparrow}_{\Delta_T}$ and $\widetilde{\epsilon}^{\uparrow}_p$ become infinite simultaneously in the limit $\Delta_T\to 0$ $(x_T\to 0)$ due to the existence of $1-[\det z_{12}]_T^{1/2}$ in the denominator. Then mathematical handling for such a problem is too difficult and therefore we can not easily get a formula for T_c^{II} in an analytical way as we did in Case I. Let us denote T_c for Case II as T_c^{II}. At $T\simeq T_c^{\mathrm{II}}$, Δ_T^{II} almost vanishes and $1-[\det z_{12}]_T^{1/2}\to 2\pi N(0)\hbar\omega_Dx_T$. Using the $\mathcal{F}^{\uparrow}_{\Delta_T}$ expressed as (3.9), we reach to the following asymptotic forms: $\mathcal{F}^{\uparrow}_{\Delta_T}\to -\,(4\pi N(0))^{-1}N(0)V\mathrm{arcsinh}\,(\hbar\omega_D/\Delta_T)$ and $(\mathcal{F}^{\uparrow}_{+\varepsilon_pT}+\mathcal{F}^{\uparrow}_{-\varepsilon_pT})/2\to(4\pi N(0))^{-1}\varepsilon_p/\Delta_T$. The QP energy $\widetilde{\epsilon}^{\uparrow}_p(=\{(\mathcal{F}^{\uparrow}_{+\varepsilon_pT}+\mathcal{F}^{\uparrow}_{-\varepsilon_pT})^2/4+\mathcal{F}^{\uparrow 2}_{\Delta_T}\}^{1/2})$ is approximately calculated as $\widetilde{\varepsilon}^{\uparrow\mathrm{II}}_p\simeq(4\pi N(0))^{-1}\varepsilon_p/\Delta_T$ $(0<\Delta_T\ll 1)$. Here we discard the contribution from $\mathcal{F}^{\uparrow}_{\Delta_T}$ comparing with the contribution from $(\mathcal{F}^{\uparrow}_{+\varepsilon_pT}+\mathcal{F}^{\uparrow}_{-\varepsilon_pT})/2$. Returning again to the original form of the BCS gap equation but with another modified QP energy $\widetilde{\epsilon}^{\uparrow\mathrm{II}}_p$, the thermal gap equation is obtained as $1=V/2\sum_p\left(1-2\widetilde{w}^{\uparrow}_p\right)/\widetilde{\epsilon}^{\uparrow\mathrm{II}}_p$ which also leads to the integral form

$$1=N(0)V\int_0^{\hbar\omega_D}d\varepsilon\frac{4\pi N(0)\Delta_T^{\mathrm{II}}}{\varepsilon}\tanh\left(\frac{\varepsilon}{2k_BT^{\mathrm{II}}\cdot 4\pi N(0)\Delta_T^{\mathrm{II}}}\right). \tag{3.28}$$

We introduce the dimensionless variable $y_T^{\mathrm{II}}\equiv\varepsilon/\left(2k_BT\cdot 4\pi N(0)\Delta_T^{\mathrm{II}}\right)$ and its upper-value $y_{T^{\mathrm{II}}}^{\mathrm{II}}\equiv\hbar\omega_D/\left(2k_BT^{\mathrm{II}}\cdot 4\pi N(0)\Delta_T^{\mathrm{II}}\right)$. Integrating (3.28) by parts $(\theta_D/k_B\widetilde{T}^{\mathrm{II}}\equiv e^C/2\pi\cdot\hbar\omega_D/k_BT^{\mathrm{II}})$, we have

$$\frac{1}{4N(0)V}\frac{1}{\pi N(0)\Delta_T^{\mathrm{II}}}\simeq\ln y_{T^{\mathrm{II}}}^{\mathrm{II}}+\ln\left(\frac{4e^C}{\pi}\right)=\ln\left(\frac{1}{\pi N(0)\Delta_T^{\mathrm{II}}}\frac{\theta_D}{\widetilde{T}^{\mathrm{II}}}\right), \tag{3.29}$$

which reads

$$\Delta_T^{\mathrm{II}}=\frac{\theta_D}{\pi N(0)\widetilde{T}^{\mathrm{II}}}\exp\left\{-\frac{1}{4N(0)V\pi N(0)\Delta_T^{\mathrm{II}}}\right\}\approx\frac{\theta_D}{\pi N(0)\widetilde{T}^{\mathrm{II}}}\left(1-\frac{1}{4N(0)V\pi N(0)\Delta_T^{\mathrm{II}}}\right). \tag{3.30}$$

From (3.30) we obtain a quadratic equation for Δ_T^{II} very near T_c and then we have a solution

$$\Delta_T^{\mathrm{II}}=\frac{1}{\pi N(0)}\frac{\theta_D}{\widetilde{T}^{\mathrm{II}}}-\frac{1}{4N(0)V\pi N(0)}, \tag{3.31}$$

in which at $T^{\mathrm{II}}=T_c^{\mathrm{II}}$, the Δ_T^{II} vanishes. Finally we can determine the critical temperature T_c^{II} for Case II as

$$T_c^{\mathrm{II}}=\frac{2e^C}{\pi}\theta_DN(0)V=1.334\theta_DN(0)V. \tag{3.32}$$

The simple formula (3.32) gives a *high* critical temperature, e.g., $T_c^{\rm II}=198$K for $N(0)V=0.25$ and $\theta_D=700$K. Finally $\Delta_T^{\rm II}$ near $T_c^{\rm II}$ can be approximately obtained as

$$\Delta_T^{\rm II}\approx-\frac{e^C}{2\pi}\frac{1}{\pi N(0)}\frac{\theta_D}{T_c^{\rm II}}\frac{T-T_c^{\rm II}}{T_c^{\rm II}},\tag{3.33}$$

which is linearly dependent on T–$T_c^{\rm II}$. It is very interesting that we could find such a dependence of $\Delta_T^{\rm II}$, comparing with the usual dependence $\sqrt{T–T_c^{\rm II}}$ of $\Delta_T^{\rm II}$. The numerical results for $N(0)V=0.25$ and θ_D=700K, obtained from (3.23), (3.24) and (3.32), are illustrated in Fig.1 below:



Figure 1: Critical Temperature T_c: **1**. $T_c^{\rm I}$=72.87K; **2**. T_c=14.49K; **3**. $T_c^{\rm II}$=198K

In the intermediate temperature region, substituting (A.16) and (A.17) into (3.16), we have

$$\left[\left(\frac{e^C}{\pi}\frac{1}{1\pm[\det z_{12}]_T^{\frac{1}{2}}}\right)^2 x_T^2\left\{N(0)V\mathrm{arcsinh}\left(\frac{1}{x_T}\right)\right\}^2-\frac{2\pi^2}{21\zeta(3)}\ln\left(\frac{e^C}{\pi}\frac{1}{1\pm[\det z_{12}]_T^{\frac{1}{2}}}\frac{\theta_D}{T}\right)\left(\frac{T}{\theta_D}\right)^2\right]$$
$$\times\left\{N(0)V\mathrm{arcsinh}\left(\frac{1}{x_T}\right)-\left(1-[\det z_{12}]_T^{\frac{1}{2}}\right)\right\}=\frac{2}{3}\left(\frac{e^C}{\pi}\frac{1}{1\pm[\det z_{12}]_T^{\frac{1}{2}}}\right)^2 x_T^2[\det z_{12}]_T^{\frac{1}{2}}.\tag{3.34}$$

A solution for equation (3.34) is classified into the following two cases:

Case I:The R.H.S. of (3.34) is approximated to be zero since $0<x_T\ll 1$, from which we obtain an equation to determine Δ_T for a given T as

$$\frac{e^C}{\pi}\frac{N(0)V}{1+[\det z_{12}]_T^{\frac{1}{2}}}x_T\mathrm{arcsinh}\left(\frac{2}{x_T}\right)=-\sqrt{\frac{2\pi^2}{21\zeta(3)}}\sqrt{\ln\left(\frac{e^C}{\pi}\frac{1}{1+[\det z_{12}]_T^{\frac{1}{2}}}\frac{\theta_D}{T}\right)}\left(\frac{T}{\theta_D}\right).\tag{3.35}$$

Using the relation and the approximations

$$\mathrm{arcsinh}\left(\frac{1}{x}\right)=\ln\left(\frac{1}{x}+\sqrt{1+\frac{1}{x^2}}\right),\ \ \frac{1}{x}+\sqrt{1+\frac{1}{x^2}}\approx\frac{1}{x}+1+\frac{1}{2}\frac{1}{x^2},\tag{3.36}$$

and $e^x\approx 1+x+\frac{1}{2}x^2$, for $[\det z_{12}]_T^{1/2}\approx 0.3$ we have

$$x_T+\frac{1}{2}=-\frac{\sqrt{0.782}}{0.436N(0)V}\sqrt{\ln\left(0.436\frac{\theta_D}{T}\right)}\left(\frac{T}{\theta_D}\right)x_T+\frac{1}{2}\frac{0.782}{\{0.436N(0)V\}^2}\ln\left(0.436\frac{\theta_D}{T}\right)\left(\frac{T}{\theta_D}\right)^2,\tag{3.37}$$

from which, finally we have a very simple solution for x_T ($N(0)V=0.25$ and $\theta_D=700$K) as

$$x_T=-\frac{1}{2}\left\{1-\frac{\sqrt{0.782}}{0.436{\times}0.25}\sqrt{\ln(0.436)-\ln\left(\frac{T}{700}\right)}\left(\frac{T}{700}\right)\right\}.\tag{3.38}$$

Case II:Using $1–[\det z_{12}]_T^{1/2} \approx 2\pi N(0)\hbar\omega_D x_T$, from (3.34) we have

$$\left[(0.283)^2\{N(0)V\}^2 \operatorname{arcsinh}^2\!\left(\frac{2}{x_T}\right) - 0.782\{\pi N(0)\hbar\omega_D\}^2\left[\ln\!\left(\frac{0.283}{2\pi N(0)\hbar\omega_D}\right) - \ln\!\left(\frac{T}{\theta_D}\right) + \operatorname{arcsinh}\!\left(\frac{2}{x_T}\right)\right]\left(\frac{T}{\theta_D}\right)^2\right] \\ \times\left\{N(0)V \operatorname{arcsinh}\!\left(\frac{2}{x_T}\right) - \left(1 - [\det z_{12}]_T^{\frac{1}{2}}\right)\right\} = -\frac{2}{3}(0.283)^2[\det z_{12}]_T^{\frac{1}{2}}. \tag{3.39}$$

Expanding (3.39) with respect to $\operatorname{arcsinh}(2/x_T)$ and neglecting a constant term which is very small for $[\det z_{12}]_T^{1/2} \approx 0.3$ and for ($N(0)V = 0.25$, $N(0)\hbar\omega_D = 0.01$ and $\theta_D = 700$K), (3.39) becomes to be a quadratic equation for $\operatorname{arcsinh}(2/x_T)$. Finally we have the following solution for x_T:

$$x_T = 2/\sinh\left[\frac{1}{2}\left\{\frac{1–0.3}{0.25} + \frac{0.782\{\pi\times 0.01\}^2}{(0.283)^2(0.25)^2}\left(\frac{T}{700}\right)^2\right\}\right. \\ \left. + \sqrt{\frac{1}{4}\left\{\frac{1–0.3}{0.25} - \frac{0.782\{\pi\times 0.01\}^2}{(0.283)^2(0.25)^2}\left(\frac{T}{700}\right)^2\right\}^2 + \frac{0.782\{\pi\times 0.01\}^2}{(0.283)^2(0.25)^3}\left[0.25\left\{\ln\!\left(\frac{0.283}{2\pi\times 0.01}\right) - \ln\!\left(\frac{T}{700}\right)\right\}\right]\left(\frac{T}{700}\right)^2}\,\right]. \tag{3.40}$$

We draw below the numerical results of the solutions for Case I and Case II.



Figure 2: Temperature dependence of the gap, Case I for $[\det z_{12}]_T^{1/2} = 0.3$, $N(0)V = 0.25$ and $\theta_D = 700$K.



Figure 3: Temperature dependence of the gap, Case II for $[\det z_{12}]_T^{1/2} = 0.3$, $N(0)V = 0.25$, $N(0)\hbar\omega = 0.01$ and $\theta_D = 700$K.

The formula for Case I gives a *high* T_c^{I}, e.g., T_c^{I}=72.87K for parameters $N(0)V$=0.25 and θ_D=700K. This is in contrast with T_c of the usual HB formula giving T_c=14.49K for the same values, $N(0)V = 0.25$ and $\theta_D = 700$. The formula for Case II gives also a *very high* T_c^{II}, e.g., T_c^{II}=198K for the same values of the parameters. They are illustrated together in Fig.1. The temperature dependence of gap near T=0 and T_c becomes more complicated than that of the HB and Abrikosov's descriptions [22, 23]. At intermediate temperature, as shown in Figs. 2 and 3, we have got the solutions of Δ_T for Cases I and II. We assume $[\det z_{12}]_T^{1/2} \approx 0.3$ (Cases I and II) and $N(0)\hbar\omega_D$=0.01 (Case II) to acquire real solutions. Anyway we could obtain really the solutions x_T=0.030 (Case I) and 0.243 (Case II) for T=80K. The former has a negative gap below 72K. To our great interest, that value is almost equal to the T_c given by (3.23). It, however, recovers a positive and small gap. Further it increases as temperature rises up to around 190K but shows vividly a decreasing tendency beyond around 200K. In this sense the former is considerably good solution. On the contrary, the latter solution naturally decreases to 0.198 as temperature rises up to around 2000K but never vanishes. This means the latter solution has no tendency approaching the T_c given by (3.32). Much improvement of the above results should be possible if the original equation (3.34) can be solved more accurately.

4 Summary and further perspectives

In the present paper we have made an attempt at a Res-MF theoretical description of the thermal behavior of the two-gap SC. To show the predominance of the Res-HBT for superconducting fermion systems with large quantum fluctuations over the usual BCS and Eliashberg's theories, we have applied the Res-HBT to the naive BCS Hamiltonian of singlet-pairing. We have derived *gap equations* within the framework of Res-HBA. From the Res-FB operators $\mathcal{F}_1$ and $\mathcal{F}_2$ with *equal-gaps*, we have found the diagonalization condition for them, which is essentially the same form as that of the BCS theory. It leads to the self-consistent Res-HB gap equation, from which we could derive the present gap. Here we have concentrated on the derivation of the *Thermal Gap Equation* with the use of the *thermal* Res-HBA. From the *thermal* Res-FB operators $\mathcal{F}_{1T}$ and $\mathcal{F}_{2T}$ with *equal-gaps* we also have found the diagonalization condition, which is just the same form of the condition used at $T=0$. This reads the self-consistent Res-HB *Thermal Gap Equation* and makes possible the derivation of the present new formulas to determine the T_c and the gaps near $T=0$ and T_c.

For *unequal two-gaps*, it is also possible to realize the above diagonalization condition for Res-FB operators $\mathcal{F}_{rp}$ $(r=1,2)$. Transforming by a unitary matrix $\widehat{g}_{rp}$, $\mathcal{F}_{rp}$ is easily diagonalized. Noticing the same correspondence as the correspondence in (3.11), $\cos\theta_{rp}\Rightarrow\cos\widehat{\theta}_{rp}$ and $\sin\theta_{rp}\Rightarrow\sin\widehat{\theta}_{rp}$, we assume each diagonalization condition (3.12) holds even in this case. Then we obtain coupled equations through a function of Δ_{1T} and Δ_{2T} expressed as

$$1=\frac{\dfrac{\varepsilon_p^2}{(\varepsilon_p^2+\Delta_{rT}^2)^{3/2}}\left(-\dfrac{2\mathcal{F}^{\uparrow}_{r\Delta_T}}{\Delta_{rT}}\right)\left(1-2\widetilde{w}^{\uparrow}_{rp}\right)}{\dfrac{\varepsilon_p}{(\varepsilon_p^2+\Delta_{rT}^2)^{3/2}}\left(\mathcal{F}^{\uparrow}_{+r\varepsilon p}+\mathcal{F}^{\uparrow}_{-r\varepsilon p}\right)\left(1-2\widetilde{w}^{\uparrow}_{rp}\right)},\quad \widetilde{w}^{\uparrow}_{rp}=\frac{1}{1+e^{\beta\widetilde{\epsilon}_{rp}}},\quad \Delta_{rT_c}=0, \tag{4.1}$$

which reduces to equation in R.H.S. of (3.11) if $\Delta_{1T}=\Delta_{2T}$. The quantities $\mathcal{F}^{\uparrow}_{r,\Delta_T}$ and $\mathcal{F}^{\uparrow}_{r,\pm\varepsilon p}$ are given by the equations similar to (5.9) in I but with more complicated forms of Δ_{1T} and Δ_{2T}. For the time being, as was done in the previous section we here also use the function $(\varepsilon_p^2+\Delta_{rT}^2)^{3/2}$ by which we divide numerator and denominator, respectively, in (4.1). After equating the numerator to the denominator and using the relation $1-2\widetilde{w}^{\uparrow}_{rp}=\tanh(\widetilde{\epsilon}_r/2k_BT)$, we sum up over p, namely integrate both sides of the equation over ε, to achieve the optimized conditions. Thus we obtain Res-HB coupled *Thermal Gap Equations* and reach our ultimate goal of computing thermal two-gaps. Along such a strategy and method, we will make a numerical analysis to demonstrate the thermal behavior of two-gaps.

To describe a superconducting fermion system with N single-particle states and to approach such the problems, we must provide a rigorous thermal Res-HBA and MF approximation. We have an expression for a partition function as $\mathrm{Tr}(e^{-\beta H})=2^{N-1}\int\langle g|e^{-\beta H}|g\rangle dg$ in an $SO(2N)$ CS rep $|g\rangle$ [19] where $\int dg$ is the group integration on group $SO(2N)$. Following Fukutome [13], introducing the projection operator P onto the Res-HB subspace, the partition function in the Res-HB subspace is computed as $\mathrm{Tr}(Pe^{-\beta H})$. This is a purely theoretical object and can be calculated within the Res-HB subspace, e.g, by using the Laplace transform of $e^{-\beta H}$ and the projection method. The result leads to an infinite matrix continued fraction, a concrete computation for which, however, is very difficult. On the other hand, a calculation of the Res-HB free energy is carried out in the same way as usual. This is made parallel to the usual thermal BCS theory, which will be given elsewhere in a separate paper in a near future.

Appendix

A Calculations of $\sum_p A_p, \sum_p B_p$ and $\sum_p C_p$ at zero and intermediate temperature

First, equation (3.16) is shown to reduce to the Res-HB gap equation (4.10) in I as $T \to 0$. Using a variable $\varepsilon = \xi \Delta_T$ instead of ε, summations $\sum_p A_p, \sum_p B_p$ and $\sum_p C_p$ near $T=0$ are computed as follows:

$$\left.\begin{aligned}
\frac{\sum_p A_p}{2N(0)} &= \operatorname{arcsinh}\left(\frac{1}{x_T}\right) - \frac{1}{\sqrt{1+x_T^2}} + A(T), \quad A(T) = -T^{\mathrm{I(II)}}_{\frac{3}{2}} + \cdots , \\
\frac{\Delta_T \sum_p B_p}{2N(0)} &= \arctan\left(\frac{1}{x_T}\right) + B(T), \quad B(T) = -T^{\mathrm{I(II)}}_{\frac{1}{2}} + \frac{1}{2} T^{\mathrm{I(II)}}_{\frac{3}{2}} - \cdots , \\
\frac{\Delta_T^2 \sum_p C_p}{2N(0)} &= \frac{1}{\sqrt{1+x_T^2}} + C(T), \quad C(T) = -T^{\mathrm{I(II)}}_{\frac{1}{2}} + T^{\mathrm{I(II)}}_{\frac{3}{2}} - \cdots,
\end{aligned}\right\} \tag{A.1}$$

detailed calculation of which is given below. With the use of the relations (3.9) and $\Delta_T/\varepsilon_p = -2\mathcal{F}^{\uparrow}_{\Delta_T}/(\mathcal{F}^{\uparrow}_{+\varepsilon_p T} + \mathcal{F}^{\uparrow}_{-\varepsilon_p T})$ which lead to $\widetilde{\varepsilon} = \sqrt{\xi^2+1}\hbar\omega_D \widetilde{\Delta}^{\mathrm{I(II)}}_T$, $\sum_p A_p$ in (3.15) is converted to

$$\frac{\sum_p A_p}{2N(0)} = \int_0^{\frac{1}{x}} d\xi \frac{\xi^2}{(\xi^2+1)^{\frac{3}{2}}} - 2\int_0^{\frac{1}{x}} d\xi \frac{\xi^2}{(\xi^2+1)^{\frac{3}{2}}} \frac{1}{1+e^{\mathbf{d}^{\mathrm{I(II)}}\sqrt{\xi^2+1}}}, \ \mathbf{d}^{\mathrm{I(II)}} \equiv \frac{\hbar\omega_D}{k_B T} \widetilde{\Delta}^{\mathrm{I(II)}}_T, \tag{A.2}$$

where $\mathbf{d}^{\mathrm{I(II)}} \gg 1$ for large $\hbar\omega_D$ and for Case I and Case II, $\widetilde{\Delta}^{\mathrm{I(II)}}_T$ is defined as

$$\widetilde{\Delta}^{\mathrm{I(II)}}_T \equiv \frac{1}{2} x_T \left\{ N(0)V \cdot \operatorname{arcsinh}\left(\frac{1}{x_T}\right) + (-)[\det z_{12}]^{\frac{1}{2}} \right\} \cdot \frac{1}{1 + (-)[\det z_{12}]^{\frac{1}{2}}}. \tag{A.3}$$

Introducing a new variable $y = \sqrt{\xi^2+1}$, (A.2) is integrated partly and approximated to be

$$\frac{\sum_p A_p}{2N(0)} \simeq \operatorname{arcsinh}\left(\frac{1}{x_T}\right) - \frac{1}{\sqrt{1+x_T^2}} - 2\int_1^{\infty} dy \frac{1}{\sqrt{y^2-1}} e^{-\mathbf{d}^{\mathrm{I(II)}}y} + 2\int_1^{\infty} dy \frac{1}{y^2} \frac{1}{\sqrt{y^2-1}} e^{-\mathbf{d}^{\mathrm{I(II)}}y}. \tag{A.4}$$

Similarly, we get approximate formulas for $\sum_p B_p$ and $\sum_p C_p$ as

$$\frac{\Delta_T \sum_p B_p}{2N(0)} \simeq \arctan\left(\frac{1}{x_T}\right) - 2\int_1^{\infty} dy \frac{1}{y} \frac{1}{\sqrt{y^2-1}} \ e^{-\mathbf{d}^{\mathrm{I(II)}}y}, \tag{A.5}$$

$$\frac{\Delta_T^2 \sum_p C_p}{2N(0)} \simeq \frac{1}{\sqrt{1+x_T^2}} - 2\int_1^{\infty} dy \frac{1}{y^2} \frac{1}{\sqrt{y^2-1}} \ e^{-\mathbf{d}^{\mathrm{I(II)}}y}. \tag{A.6}$$

To carry out integral calculations in (A.4) $\sim$ (A.6), it is convenient to use an integral representation of Bessel function [26]. The Bessel function of order ν is represented as

$$K_\nu(z) = \frac{\sqrt{\pi}\left(\frac{z}{2}\right)^\nu}{\Gamma\left(\nu + \frac{1}{2}\right)} \int_1^{\infty} dy (y^2-1)^{\nu-\frac{1}{2}} e^{-zy}. \tag{A.7}$$

Then, the integral form of the Bessel function of order 0 and its exact result are given by

$$K_0(z) = \frac{\sqrt{\pi}}{\Gamma\left(\frac{1}{2}\right)} \int_1^{\infty} dy \frac{1}{\sqrt{y^2-1}} \ e^{-zy} = \sqrt{\frac{\pi}{2z}} e^{-z}, \ (\Gamma\left(\tfrac{1}{2}\right) = \sqrt{\pi}). \tag{A.8}$$

Using (A.8), integral calculations of (A.4) $\sim$ (A.6) are made in the following ways:

$$\begin{aligned}
\int_1^{\infty} dy \frac{1}{y} \frac{1}{\sqrt{y^2-1}} \ e^{-\mathbf{d}y} &= \int_1^{\infty} dy \frac{1}{\sqrt{y^2-1}} \int_{\mathbf{d}}^{\infty} dz e^{-zy} = \int_{\mathbf{d}}^{\infty} dz K_0(z) \\
&= \sqrt{\frac{\pi}{2}} \left(\mathbf{d}^{-\frac{1}{2}} - \frac{1}{2}\mathbf{d}^{-\frac{3}{2}} + \frac{3}{4}\mathbf{d}^{-\frac{5}{2}} - \cdots \right) e^{-\mathbf{d}},
\end{aligned} \tag{A.9}$$

$$
\begin{aligned}
\int_1^\infty dy \frac{1}{y^2} \frac{1}{\sqrt{y^2-1}}\ e^{-\mathbf{d}y} &= \int_1^\infty dy \frac{1}{y} \frac{1}{\sqrt{y^2-1}} \int_{\mathbf{d}}^\infty dw e^{-wy} = \int_{\mathbf{d}}^\infty dw \int_w^\infty dz K_0(z) \\
&= \sqrt{\frac{\pi}{2}} \left(\mathbf{d}^{-\frac{1}{2}} - \mathbf{d}^{-\frac{3}{2}} + \frac{9}{4} \mathbf{d}^{-\frac{5}{2}} - \cdots \right) e^{-\mathbf{d}}.
\end{aligned}
\tag{A.10}
$$

As a result, we obtain the approximation for $A(T)$, $B(T)$ and $C(T)$ near $T=0$ as

$$
A(T) \simeq 0, \quad B(T) = C(T) \simeq -T_{\frac{1}{2}}^{\mathrm{I(II)}}.
\tag{A.11}
$$

Further in the above near $T=0$ we make the following approximations:

$$
\left.
\begin{aligned}
\arctan\left(\frac{1}{x_T}\right) \simeq \frac{\pi}{2} - x_T, \quad \frac{1}{\sqrt{1+x_T^2}} \simeq 1 - x_0 x_T, \quad (0 < x_0 \ll 1) \\
\operatorname{arcsinh}\left(\frac{1}{x_T}\right) \simeq \operatorname{arcsinh}\left(\frac{1}{x_0}\right) - \frac{1}{x_0}(x_T - x_0), \quad [\det z_{12}]_T^{\frac{1}{2}} \simeq [\det z_{12}]_{T=0}^{\frac{1}{2}}.
\end{aligned}
\right\}
\tag{A.12}
$$

Next, let us introduce a new variable y by $\varepsilon = 4(1 \pm [\det z_{12}]_T^{1/2}) k_B T y$ and quantities $\widetilde{x}_T = \widetilde{\Delta}_T / \hbar\omega_D$ and $y_T^{(\pm)} = \sqrt{\varepsilon^2 + \widetilde{\Delta}_T^2} \{4(1 \pm [\det z_{12}]_T^{1/2})\}^{-1} \hbar\omega_D / k_B T$ where $\widetilde{\Delta}_T \equiv \Delta_T N(0) V \operatorname{arcsinh}(\hbar\omega_D / \Delta_T)$. In intermediate temperature region the modified QP energy $\widetilde{\varepsilon}$ is approximated as $\widetilde{\varepsilon}^{(\pm)} = \sqrt{\varepsilon^2 + \widetilde{\Delta}_T^2} \{2(1 \pm [\det z_{12}]_T^{1/2})\}^{-1}$. If $\varepsilon \gg \Delta_T$, $\sum_p A_p$, $\sum_p B_p$ and $\sum_p C_p$ in (3.15) are recast to the following integrals up to $\widetilde{\Delta}_T$:

$$
\begin{aligned}
\frac{\sum_p A_p}{2N(0)} &\simeq \int_0^{\hbar\omega_D} d\varepsilon \frac{1}{\sqrt{\varepsilon^2 + \widetilde{\Delta}_T^2}} \tanh\left(\frac{\widetilde{\varepsilon}^{(\pm)}}{2k_B T}\right) - \int_0^{\hbar\omega_D} d\varepsilon \frac{\widetilde{\Delta}_T^2}{\varepsilon^2 \sqrt{\varepsilon^2 + \widetilde{\Delta}_T^2}} \tanh\left(\frac{\widetilde{\varepsilon}^{(\pm)}}{2k_B T}\right) \\
&= \int_0^{y_T^{(\pm)}} dy \left\{ 1 - \frac{\left(y_T^{(\pm)} \widetilde{x}_T\right)^2}{y^2} \right\} \frac{1}{\sqrt{y^2 + \left(y_T^{(\pm)} \widetilde{x}_T\right)^2}} \tanh\left[\sqrt{y^2 + \left(y_T^{(\pm)} \widetilde{x}_T\right)^2}\right],
\end{aligned}
\tag{A.13}
$$

$$
\frac{\widetilde{\Delta}_T \sum_p B_p}{N(0)} \simeq \int_{-\hbar\omega_D}^{\hbar\omega_D} d\varepsilon \frac{\widetilde{\Delta}_T}{\varepsilon \sqrt{\varepsilon^2 + \widetilde{\Delta}_T^2}} \tanh\left(\frac{\widetilde{\varepsilon}^{(\pm)}}{2k_B T}\right) - \frac{1}{2} \int_{-\hbar\omega_D}^{\hbar\omega_D} d\varepsilon \frac{\widetilde{\Delta}_T^3}{\varepsilon^3 \sqrt{\varepsilon^2 + \widetilde{\Delta}_T^2}} \tanh\left(\frac{\widetilde{\varepsilon}^{(\pm)}}{2k_B T}\right) = 0,
\tag{A.14}
$$

$$
\begin{aligned}
\frac{\widetilde{\Delta}_T^2 \sum_p C_p}{2N(0)} &\simeq \int_0^{\hbar\omega_D} d\varepsilon \frac{\widetilde{\Delta}_T^2}{\varepsilon^2 \sqrt{\varepsilon^2 + \widetilde{\Delta}_T^2}} \tanh\left(\frac{\widetilde{\varepsilon}^{(\pm)}}{2k_B T}\right) - \int_0^{\hbar\omega_D} d\varepsilon \frac{\widetilde{\Delta}_T^4}{\varepsilon^4 \sqrt{\varepsilon^2 + \widetilde{\Delta}_T^2}} \tanh\left(\frac{\widetilde{\varepsilon}^{(\pm)}}{2k_B T}\right) \\
&= \left(y_T^{(\pm)} \widetilde{x}_T\right)^2 \int_0^{y_T^{\mathrm{II}}} dy \left\{ 1 - \frac{\left(y_T^{(\pm)} \widetilde{x}_T\right)^2}{y^2} \right\} \frac{1}{y^2 \sqrt{y^2 + \left(y_T^{(\pm)} \widetilde{x}_T\right)^2}} \tanh\left[\sqrt{y^2 + \left(y_T^{(\pm)} \widetilde{x}_T\right)^2}\right].
\end{aligned}
\tag{A.15}
$$

Further equations (A.13) and (A.15) are approximately computed, respectively, as

$$
\left.
\begin{aligned}
\frac{\sum_p A_p}{2N(0)} &= \int_0^{\hbar\omega_D} d\varepsilon \frac{\varepsilon^2}{\left(\varepsilon^2 + \widetilde{\Delta}_T^2\right)^{\frac{3}{2}}} \tanh\left(\frac{\widetilde{\varepsilon}^{(\pm)}}{2k_B T}\right) \simeq \ln\left(\frac{4e^C}{\pi} y_T^{(\pm)}\right) - \frac{21}{2\pi^2} \zeta(3) \left(y_T^{(\pm)} \widetilde{x}_T\right)^2, \\
\frac{\widetilde{\Delta}_T^2 \sum_p C_p}{2N(0)} &= \int_0^{\hbar\omega_D} d\varepsilon \frac{1}{\left(\varepsilon^2 + \widetilde{\Delta}_T^2\right)^{\frac{3}{2}}} \tanh\left(\frac{\widetilde{\varepsilon}^{(\pm)}}{2k_B T}\right) \simeq \frac{7}{\pi^2} \zeta(3) \left(y_T^{\mathrm{II}} x_T\right)^2.
\end{aligned}
\right\}
\tag{A.16}
$$

Taking only a leading term, finally terms A_p and C_p in (A.16) are approximated to be

$$\left.\begin{aligned} &\frac{\sum_p A_p}{2N(0)} \simeq \ln\left(\frac{e^C}{\pi}\frac{1}{1\pm[\det z_{12}]_T^{\frac{1}{2}}}\frac{\theta_D}{T}\right) - \alpha_T^{(\pm)}, \;\; \frac{\widetilde{\Delta}_T^2 \sum_p C_p}{2N(0)} \simeq \frac{2}{3}\alpha_T^{(\pm)}, \\ &\alpha_T^{(\pm)} \equiv \frac{21\zeta(3)}{2\pi^2}\left(\frac{e^C}{\pi}\frac{1}{1\pm[\det z_{12}]_T^{\frac{1}{2}}}\right)^2\left(\frac{\widetilde{\Delta}_T}{k_B T}\right)^2. \end{aligned}\right\} \tag{A.17}$$

To derive (A.16) we give a integral formula

$$\int_0^\infty dy \left\{\frac{1}{y^3}\tanh y - \frac{1}{y^2}\mathrm{sech}^2 y\right\} = \frac{7}{\pi^2}\zeta(3), \tag{A.18}$$

which can be derived by using the famous mathematical formulas [24]

$$\frac{1}{y}\tanh y = 8\sum_{m=1}^\infty \frac{1}{(2m-1)^2\pi^2+4y^2}, \;\; \frac{1}{y^3}\tanh y - \frac{1}{y^2}\mathrm{sech}^2 y = 64\sum_{m=1}^\infty \frac{1}{\{(2m-1)^2\pi^2+4y^2\}^2}. \tag{A.19}$$

Adopting a new integral variable $y = (2m-1)\pi/2 \cdot \tan\theta$, an integral of the second formula in (A.19) is easily carried out for $\hbar\omega_D \gg 1$ as

$$64\int_0^{y_T^{\mathrm{II}}\to\infty} dy \sum_{m=1}^\infty \frac{1}{\{(2m-1)^2\pi^2+4y^2\}^2} = 32\sum_{m=1}^\infty \frac{1}{(2m-1)^3\pi^3}\int_0^{\frac{\pi}{2}} d\theta \frac{1}{1+\tan^2\theta} = \frac{7}{\pi^2}\zeta(3), \tag{A.20}$$

where we have used $\sum_{m=1}^\infty (2m-1)^{-3} = (7/8)\cdot\zeta(3)$, and $\zeta(3) = \pi^3/25.79436$ [24].

To get a finite value of $\sum_p A_p$, expanding (A.13) around $y_T^{(\pm)}\widetilde{x}_T$, (A.13) is boldly approximated as

$$\frac{\sum_p A_p}{2N(0)} \simeq \int_0^{y_T^{(\pm)}} dy \frac{1}{y}\tanh y - \frac{3}{2}\left(y_T^{(\pm)}\widetilde{x}_T\right)^2 \int_0^{y_T^{(\pm)}} dy \left\{\frac{1}{y^3}\tanh y - \frac{1}{y^2}\mathrm{sech}^2 y\right\}. \tag{A.21}$$

In a similar way we also get a roughly approximated integral form for (A.15) as

$$\frac{\widetilde{\Delta}_T^2 \sum_p C_p}{2N(0)} \simeq \left(y_T^{(\pm)}\widetilde{x}_T\right)^2 \int_0^{y_T^{(\pm)}} dy \left\{\frac{1}{y^3}\tanh y - \frac{1}{y^2}\mathrm{sech}^2 y\right\}. \tag{A.22}$$

Integrations of (A.21) and (A.22) are easily made by using the integration formula (A.18) if we take the upper-value $y_T^{(\pm)}$ to be infinite.

Acknowledgements

S. N. would like to express his sincere thanks to Professor Manuel Fiolhais for kind and warm hospitality extended to him at the Centro de Física Computacional, Universidade de Coimbra, Portugal. This work was supported by FCT (Portugal) under the project CERN/FP/83505/2008.

References

[1] J. Nagamatsu, N. Nakagawa, T. Muranaka, Y. Zenitani and J. Akimitsu, Nature **410** (2001), 63-64

[2] J. Bardeen, L.N. Cooper and J.R. Schrieffer, Phys. Rev. **108** (1957), 1175-1204

[3] N.N. Bogoliubov, Usp. Fiz. Nauk **67** (1959), 549-580 [Soviet Phys. Uspekhi **67** (1959), 236-254]

[4] N.N. Bogoliubov, V.V. Tolmachev, D.V. Shirkov, *A New Method in the Theory of Superconductivity* (Consultants Bureau, New York, 1959)

[5] G.M. Eliashberg, Zh. Exp. Teor. Fiz. **38** (1960), 966-976 [Soviet Phys. JETP **11** (1960), 696 -702]

[6] J.R. Schrieffer, *Theory of Superconductivity* (W.A. Benjamine, New York, 1964)

[7] *The Problem of High Temperature Superconductivity*, edited by V.L. Ginzburg and D.A. Kirzhnits (Consultant Bureau, New York, 1982)

[8] R.D. Parks (ed.), *Superconductivity*, 2 Vols. (Marcel Dekker, New York, 1969)

[9] Jens Kortus, I.I. Mazin, K.D. Belashchenko, V.P. Antropov and L.L. Boyer, Phys. Rev. Lett. **86** (2001), 4656-4659

[10] Amy Y. Liu, I.I. Mazin and Jens Kortus, Phys. Rev. Lett. **87** (2001), 87005-1-87005-4

[11] H. Shul, B.T. Matthias and L.R. Walker, Phys. Rev. Lett. **3** (1959), 552-554

[12] J. Kondo, Prog. Theor. Phys. **29** (1963), 1-9

[13] H. Fukutome, Prog. Theor. Phys. **80** (1988), 417-432

[14] S. Nishiyama and H. Fukutome, Prog. Theor. Phys. **85** (1991), 1211-1222

[15] S. Nishiyama and H. Fukutome, J. Phys. G: Nucl. Part. Phys. **18** (1992), 317-328;
H. Ohnishi, J. da Providência and S. Nishiyama, in *Proceedings of the 29th International Workshop on Condensed Matter Theories*, ed. H. Akai, A. Hosaka, H. Toki and F. Bary Malik, (Nova Science Publishers, Inc., USA), Condensed Matter Theories Vol. **21** (2007), 395-406

[16] S. Nishiyama, J. da Providência, C. Providência and H. Ohnishi, Adv. Studies Theor. Phys. **4** (2010), 283-303.

[17] P.G. de Gennes, *Superconductivity of metals and alloys*, (Benjamine, New York, 1966)

[18] P. Ring and P. Schuk, *The nuclear many body problem*, Texts and monographs in Physics, (Springer, Berlin, Heidelberg and New York, 1980)

[19] A.M. Perelomov, Commun. Math. Phys. **26** (1972), 222-236; Usp. Fiz. Nauk **123** (1977), 23-55 [Sov. Phys. Usp. **20** (1977), 703-720]

[20] S. Nishiyama, J. da Providência and H. Ohnishi, in *Proceedings of the 26th International Workshop on Condensed Matter Theories*, ed. M. de Llano, C. Fiolhais and J. da Providência, (Nova Science Publishers, Inc., USA), Condensed Matter Theories Vol. **18** (2003), 169-178; in *Proceedings of the 29th International Workshop on Condensed Matter Theories*, ed. H. Akai, A. Hosaka, H. Toki and F. Bary Malik, (Nova Science Publishers, Inc., USA), ibid. Vol. **21** (2007), 13-25

[21] I.M. Khalatnikov and A.A. Abrikosov, Adv. Phys. **8** (1959), 45-86

[22] A.A. Abrikosov, L.P. Gor'kov and I. Ye. Dzyaloshinskii, *Quantum Field Theoretical Methods in Statistical Physics*, (Pergamon, Oxford, 1965)

[23] A.A. Abrikosov, *Fundamentals of the Theory of Metals*, (North-Holland, Amsterdam, 1988)

[24] I.S. Gradshteyn and I.M. Ryzhik, A. Jeffrey, Editor, *Tables of Integrals, Series and Products*, Fifth Edition, Academic Press, London, 1994.

[25] G.M. Eliashberg, Zh. Exp. Teor. Fiz. **39** (1960), 1437-1441 [Soviet Phys. JETP, **12** (1961), 1000-1002]

[26] G.N. Watson, *A treatise on the theory of Bessel functions*, Second edition, Cambridge University Press, Cambridge, 1958.